



**AS**

# **Physics data and formulae**

**For use in exams from the June 2016 Series onwards**

**DATA - FUNDAMENTAL CONSTANTS AND VALUES**

<b>Quantity</b>	<b>Symbol</b>	<b>Value</b>	<b>Units</b>
speed of light in vacuo	$c$	$3.00 \times 10^8$	$\text{m s}^{-1}$
permeability of free space	$\mu_0$	$4\pi \times 10^{-7}$	$\text{H m}^{-1}$
permittivity of free space	$\epsilon_0$	$8.85 \times 10^{-12}$	$\text{F m}^{-1}$
magnitude of the charge of electron	$e$	$1.60 \times 10^{-19}$	$\text{C}$
the Planck constant	$h$	$6.63 \times 10^{-34}$	$\text{J s}$
gravitational constant	$G$	$6.67 \times 10^{-11}$	$\text{N m}^2 \text{kg}^{-2}$
the Avogadro constant	$N_A$	$6.02 \times 10^{23}$	$\text{mol}^{-1}$
molar gas constant	$R$	8.31	$\text{J K}^{-1} \text{mol}^{-1}$
the Boltzmann constant	$k$	$1.38 \times 10^{-23}$	$\text{J K}^{-1}$
the Stefan constant	$\sigma$	$5.67 \times 10^{-8}$	$\text{W m}^{-2} \text{K}^{-4}$
the Wien constant	$\alpha$	$2.90 \times 10^{-3}$	$\text{m K}$
electron rest mass (equivalent to $5.5 \times 10^{-4}$ u)	$m_e$	$9.11 \times 10^{-31}$	$\text{kg}$

electron charge/mass ratio	$\frac{e}{m_e}$	$1.76 \times 10^{11}$	C kg <sup>-1</sup>
proton rest mass (equivalent to 1.00728 u)	$m_p$	$1.67(3) \times 10^{-27}$	kg
proton charge/mass ratio	$\frac{e}{m_p}$	$9.58 \times 10^7$	C kg <sup>-1</sup>
neutron rest mass (equivalent to 1.00867 u)	$m_n$	$1.67(5) \times 10^{-27}$	kg
gravitational field strength	$g$	9.81	N kg <sup>-1</sup>
acceleration due to gravity	$g$	9.81	m s <sup>-2</sup>
atomic mass unit (1u is equivalent to 931.5 MeV)	u	$1.661 \times 10^{-27}$	kg

[Turn over]

**ALGEBRAIC EQUATION**

quadratic equation  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

**ASTRONOMICAL DATA**

Body	Mass/kg	Mean radius/m
Sun	$1.99 \times 10^{30}$	$6.96 \times 10^8$
Earth	$5.97 \times 10^{24}$	$6.37 \times 10^6$

**GEOMETRICAL EQUATIONS**

arc length  $= r\theta$

circumference of circle  $= 2\pi r$

area of circle  $= \pi r^2$

curved surface area of cylinder  $= 2\pi rh$

area of sphere  $= 4\pi r^2$

volume of sphere  $= \frac{4}{3}\pi r^3$

## Particle Physics

Class	Name	Symbol	Rest energy/MeV
photon	photon	$\gamma$	0
lepton	neutrino	$\nu_e$	0
		$\nu_\mu$	0
	electron	$e^\pm$	0.510999
	muon	$\mu^\pm$	105.659
mesons	$\pi$ meson	$\pi^\pm$	139.576
		$\pi^0$	134.972
	K meson	$K^\pm$	493.821
		$K^0$	497.762
baryons	proton	p	938.257
	neutron	n	939.551

[Turn over]

**Properties of quarks**  
**antiquarks have opposite signs**

Type	Charge	Baryon number	Strangeness
u	$+\frac{2}{3}e$	$+\frac{1}{3}$	0
d	$-\frac{1}{3}e$	$+\frac{1}{3}$	0
s	$-\frac{1}{3}e$	$+\frac{1}{3}$	-1

## Properties of Leptons

		Lepton number
Particles:	$e^-, \nu_e; \mu^-, \nu_\mu$	+ 1
Antiparticles:	$e^+, \bar{\nu}_e, \mu^+, \bar{\nu}_\mu$	- 1

## Photons and energy levels

photon energy

$$E = hf = hc / \lambda$$

photoelectricity

$$hf = \phi + E_k (\text{max})$$

energy levels

$$hf = E_1 - E_2$$

de Broglie wavelength

$$\lambda = \frac{h}{p} = \frac{h}{mv}$$

[Turn over]

## Waves

wave speed  $c = f\lambda$       period  $f = \frac{1}{T}$

first harmonic  $f = \frac{1}{2l} \sqrt{\frac{T}{\mu}}$

fringe spacing  $w = \frac{\lambda D}{s}$       diffraction grating  $d \sin \theta = n\lambda$

refractive index of a substance  $s$ ,  $n = \frac{c}{c_s}$

for two different substances of refractive indices  $n_1$  and  $n_2$ ,

law of refraction  $n_1 \sin \theta_1 = n_2 \sin \theta_2$

critical angle  $\sin \theta_c = \frac{n_2}{n_1}$  for  $n_1 > n_2$



**Mechanics****moments**

**moment =  $Fd$**

**velocity and  
acceleration**

**$v = \frac{\Delta s}{\Delta t}$**

**$a = \frac{\Delta v}{\Delta t}$**

**equations of  
motion**

**$v = u + at$**

**$s = \left( \frac{u+v}{2} \right) t$**

**$v^2 = u^2 + 2as$**

**$s = ut + \frac{at^2}{2}$**

**force**

**$F = ma$**

**force**

**$F = \frac{\Delta(mv)}{\Delta t}$**

**impulse**

**$F \Delta t = \Delta(mv)$**

**work, energy  
and power**

**$W = F s \cos \theta$**

**$E_k = \frac{1}{2} m v^2$**

**$\Delta E_p = mg \Delta h$**

**$P = \frac{\Delta W}{\Delta t}$  ,  **$P = Fv$****

**$efficiency = \frac{useful\ output\ power}{input\ power}$**

**[Turn over]**

**Materials**

density  $\rho = \frac{m}{V}$

Hooke's law  $F = k \Delta L$

Young modulus =  $\frac{\text{tensile stress}}{\text{tensile strain}}$

tensile stress =  $\frac{F}{A}$

tensile strain =  $\frac{\Delta L}{L}$

energy stored  $E = \frac{1}{2} F \Delta L$

**Electricity**

current and pd  $I = \frac{\Delta Q}{\Delta t}$      $V = \frac{W}{Q}$      $R = \frac{V}{I}$

resistivity  $\rho = \frac{RA}{L}$

resistors in series  $R_T = R_1 + R_2 + R_3 + \dots$

resistors in parallel  $\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$

power  $P = VI = I^2 R = \frac{V^2}{R}$

emf  $\varepsilon = \frac{E}{Q}$      $\varepsilon = I(R + r)$

**There are no formulae printed on this page**

**Copyright Information**

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from [www.aqa.org.uk](http://www.aqa.org.uk) after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2016 AQA and its licensors. All rights reserved.

**IB/M/Jun16/CD/7407/INS/E7**