CAMBRIDGE INTERNATIONAL EXAMINATIONS GCE Advanced Subsidiary Level and GCE Advanced Level

www.papacambridge.com MARK SCHEME for the October/November 2012 series

9700 BIOLOGY

9700/23


Paper 2 (AS Structured Questions), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.


Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2012 series for most IGCSE. GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



F	Page 3	Mark Scheme	Syllabu er
		GCE AS/A LEVEL – October/November 2012	9700 2020
	accept first on accept phonet		Syllabu, Dana er 9700 Para annbra
n	name	mitochondrion ; A mitochondria	
fı	<i>unction</i> (site c	of); <u>ATP</u> , synthesis / production / AW <u>aerobic</u> respiration link reaction Krebs cycle oxidative phosphorylation AVP R ATP energy	
n n	ame	Golgi (apparatus / body / complex) ; A dictyosome A Golgi	
fL	<i>unction</i> (site c	of); modification of protein / glycosylation / describe modification of lipid pack(ag)ing (of), protein / lipids production of (Golgi / secretory) vesicles / lyso ignore synthesis of protein (<i>incorrect name</i>) <i>lysosome function</i> = contains / storage of hydro <i>Golgi / secretory, vesicles</i> = transport, protein	somes olytic / digestive, enzymes
; n	ame	chloroplast(s) ;	
fL	<i>unction</i> (site c	of); photosynthesis light-dependent, reactions / stage (of photosyn light, absorption / AW light-independent, reactions / stage (of photosy Calvin cycle carbon fixation photophosphorylation A ATP synthesis ignore (<i>treat as neutral</i>) <i>ref. to, glucose / oxyg</i> ignore <i>chlorophyll</i> R light / dark, stage / reactions	ynthesis)
n	name	<u>rough</u> endoplasmic reticulum ; R RER or rough ER R endoplastic	
fı	<i>unction</i> (site c	of); protein / polypeptide, synthesis translation modification of protein / described (e.g. folding protein transport (to Golgi) <i>(incorrect name)</i> <i>smooth endoplasmic reticulum</i> = lipid / steroid <i>endoplasmic reticulum</i> = <i>ecf as above for RER</i>	/ cholesterol, synthesis / AW



Page 5	Mark Scheme	Syllabu & er
	GCE AS/A LEVEL – October/November 2012	9700

- 3 (a) look at quoted data to confirm qualitative statements if unclear
- cambridge.com 1 people who never smoked have the lowest percentage of deaths (due to lung cance must be comparative

for age

2 either

the younger / earlier the person starts smoking the higher the percentage of deaths or

the older / later the person starts smoking the lower the percentage of deaths (due to lung cancer);

for number of cigarettes per day

3 either

increasing / AW, the number of cigarettes smoked per day increases the percentage of deaths

or

decreasing / AW, the number of cigarettes smoked per day decreases the percentage of deaths ;

different 'start' ages for the two types of smokers

- highest percentage deaths is for those with an early start and smoke, 21-39 (cigarettes 4 per day) / the most / AW ;
- 5 greatest difference in percentage deaths occurs in those that start smoking early; ora [max 4]
- (b) (i) 1 forms carboxyhaemoglobin ;
 - reduces affinity of Hb for oxygen / Hb has higher affinity for CO than for oxygen ; 2 ignore 'picks up CO rather than oxygen', if mp3 is given then allow
 - 3 reduces quantity of oxygen transported (in blood) / AW ; **R** prevents
 - 4 damages lining of arteries ; A promotes / AW, atheroma / atherosclerosis / plaque [max 2]
 - (ii) raises, heart rate / blood pressure ; reduces diameter of arterioles ; decreases blood flow to body extremities ; increases 'stickiness' of platelets / promotes, blood clotting / thrombosis ; [max 2]

	Mark Scheme	Syllabu er					
	GCE AS/A LEVEL – October/November 2012	Syllabu 9700 Bolgi bodies or vesicles					
(iii) goblet cells							
(111)	enlarge / swell up ;	76.					
	A become bigger / dilate	10					
	R inflamed						
	produce more / excess, mucus ;						
	A lots of						
AVP ; e.g. any cellular detail such as more mitochondria / Golgi bodies or ve							
	cilia:						
	paralysis / destruction ;						
	A damages R kills <i>ignore</i> 'tar coats…'						
	no / less beating / sweeping (action) / moving mucus ;						
	R in context of moving air	[max 4]					
		[Total: 12]					
(a) (x) 4	(a) (×) 400 ;;						
• • • •	if answer incorrect or not to nearest 100 allow one mark for correct working						
	e.g. (scale bar) 19 000–21 000 divided by 50						
-	rd max one mark if a unit (e.g. μ m) is included	[2]					
(b) 1	thick(ened) / lignified, walls prevent, collapse ;						
	ignore strenghtened						
	A withstands, compression / negative pressure						
2	ignore bursting lignified (wall), prevents leakage / provides waterproofing ;						
3	cellulose, wall / lining, allows adhesion of water (molecules)	<u>ر</u> .					
0	A hydrogen bonding / hydrophilic	·/ ,					
4	(relatively) large diameter / large cross-sectional area / wide	e / large lumen :					
	hollow / empty / no contents / no cytoplasm ;						
	no end walls / continuous 'tubes' / AW ;						
5							
5 6	elongated ;						
5 6	•	ake tubes)					
5 6	elongated ; A <i>if referenced to cells or vessels</i> A cells end to end (to ma only allow mps 4–7 in terms of ease / efficiency of water mo	ovement					
5 6	elongated ; A <i>if referenced to cells or vessels</i> A <i>cells end to end (to ma only allow mps 4–7 in terms of ease / efficiency of water mo mp 4 e.g. more space allows a greater volume to flow / greater volume to flow /</i>	ovement ater volume per unit time					
5 6 7	elongated ; A <i>if referenced to cells or vessels</i> A cells end to end (to ma <i>only allow mps</i> 4–7 <i>in terms of ease / efficiency of water mo</i> <i>mp</i> 4 e.g. more space allows a greater volume to flow / greater <i>or mp</i> 5–6 e.g. <i>minimal resistance to flow, allows unimpede</i>	ovement ater volume per unit time					
5 6	elongated ; A <i>if referenced to cells or vessels</i> A <i>cells end to end (to ma only allow mps 4–7 in terms of ease / efficiency of water mo mp 4 e.g. more space allows a greater volume to flow / greater volume to flow /</i>	ovement ater volume per unit time					

	ige 7	Mark Scheme Syllabu	er
		GCE AS/A LEVEL – October/November 2012 9700	03
		Mark Scheme Syllabu. GCE AS/A LEVEL – October/November 2012 9700 water moves, down a water potential gradient / from a high(er) water potential low(er) water potential, accept ψ for water potential ; apoplast pathway, described / used in correct context ; symplast pathway, described / used in correct context ; evaporation from mesophyll cell walls ; A surface of mesophyll cells into air space(s) :	°C2.
(c)	1	water moves, down a <u>water potential</u> gradient / from a high(er) water potential	76.
	~	low(er) water potential, accept ψ for water potential ;	1
	2	apoplast pathway, described / used in correct context ;	
	3	symplast pathway, described / used in correct context ;	
	4	evaporation from mesophyll cell walls ;	
	F	A surface of mesophyll cells	
	5		
	c	must be linked to evaporation / water vapour	
	6	water vapour diffuses (out);	
	7	accept if no vapour but follows from evaporation	
	1	out / through / via <u>stoma(</u> ta) ; R 'evaporates from the stomata'	
	8	AVP ; ref. to water leaves unlignified terminals of xylem vessels	[max 5
	0	AVF, Tel. 10 water leaves unlightned terminals of Aylent vessels	linax o
			[Total: 10]
(2)		points assant mn2 may be taken from a labelled/annatated disgram	
(a)		points except mp3 may be taken from a labelled/annotated diagram ref. to, attachment / AW, to mRNA ;	
	1 2	<i>idea of</i> two codon attachment, sites / space, for six bases <i>or</i> nucleotides ;	
	2	mRNA has code for sequence of amino acids (in a polypeptide);	
	4	(ribosome) provides sites for attachment of two tRNA (molecules) ;	
	т	A implied	
	5	each tRNA has a specific amino acid / AW ;	
	6	(mRNA) codon – anticodon (tRNA), binding ;	
	-	A description in terms of complementary base pairing	
		A 'matching'	
	7	formation of peptide bonds (catalysed by peptidyl transferase);	
	8	idea of ribosome moving along mRNA one codon at a time ;	[max 4
(b)	(i)	GGC ;	[1]
	(;;)	CTA;	[4
	(11)		[1
(c)	1	amino acid coded by codon 2 changed ;	
(c)	1 2	amino acid coded by codon 2 changed ; <i>idea of</i> every subsequent <u>codon</u> changed ;	
(c)			
(c)	2	idea of every subsequent codon changed ;	
(c)	2	<i>idea of</i> every subsequent <u>codon</u> changed ; amino acids / protein sequence, up to and including codon 1 unaffected / AW	
(c)	2 3	<i>idea of</i> every subsequent <u>codon</u> changed ; amino acids / protein sequence, up to and including codon 1 unaffected / AW ora amino acid sequence from codon 2 onwards is changed ;	
(c)	2 3 4	<i>idea of</i> every subsequent <u>codon</u> changed ; amino acids / protein sequence, up to and including codon 1 unaffected / AW ora amino acid sequence from codon 2 onwards is changed ; <i>idea of</i> premature chain termination (if stop codon further on) / AW ;	
(c)	2 3 4 5	<i>idea of</i> every subsequent <u>codon</u> changed ; amino acids / protein sequence, up to and including codon 1 unaffected / AW ora amino acid sequence from codon 2 onwards is changed ; <i>idea of</i> premature chain termination (if stop codon further on) / AW ; <i>idea of</i> change in, <u>primary</u> / <u>secondary</u> / <u>tertiary</u> , structure of protein ;	
(c)	2 3 4 5	<i>idea of</i> every subsequent <u>codon</u> changed ; amino acids / protein sequence, up to and including codon 1 unaffected / AW ora amino acid sequence from codon 2 onwards is changed ; <i>idea of</i> premature chain termination (if stop codon further on) / AW ; <i>idea of</i> change in, <u>primary</u> / <u>secondary</u> / <u>tertiary</u> , structure of protein ; <i>idea of</i> protein non-functional ;	
(c)	2 3 4 5	 <i>idea of</i> every subsequent <u>codon</u> changed ; amino acids / protein sequence, up to and including codon 1 unaffected / AW ora amino acid sequence from codon 2 onwards is changed ; <i>idea of</i> premature chain termination (if stop codon further on) / AW ; <i>idea of</i> change in, <u>primary</u> / <u>secondary</u> / <u>tertiary</u>, structure of protein ; <i>idea of</i> protein non-functional ; ignore 'affect / effect' A in context of enzyme not functioning R if this point is out of context 	
(c)	2 3 4 5	 <i>idea of</i> every subsequent <u>codon</u> changed ; amino acids / protein sequence, up to and including codon 1 unaffected / AW ora amino acid sequence from codon 2 onwards is changed ; <i>idea of</i> premature chain termination (if stop codon further on) / AW ; <i>idea of</i> change in, <u>primary</u> / <u>secondary</u> / <u>tertiary</u>, structure of protein ; <i>idea of</i> protein non-functional ; ignore 'affect / effect' A in context of enzyme not functioning 	[max 3
(c)	2 3 4 5 6	 <i>idea of</i> every subsequent <u>codon</u> changed ; amino acids / protein sequence, up to and including codon 1 unaffected / AW ora amino acid sequence from codon 2 onwards is changed ; <i>idea of</i> premature chain termination (if stop codon further on) / AW ; <i>idea of</i> change in, <u>primary</u> / <u>secondary</u> / <u>tertiary</u>, structure of protein ; <i>idea of</i> protein non-functional ; ignore 'affect / effect' A in context of enzyme not functioning R if this point is out of context 	[max 3 [Total: 9

