

MARK SCHEME for the May/June 2014 series

9700 BIOLOGY

9700/42

Paper 4 (A2 Structured Questions), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

Mark scheme abbreviations:

;	separates marking points
/	alternative answers for the same point
R	reject
A	accept (for answers correctly cued by the question, or by extra guidance)
AW	alternative wording (where responses vary more than usual)
<u>underline</u>	actual word given must be used by candidate (grammatical variants accepted)
max	indicates the maximum number of marks that can be given
ora	or reverse argument
mp	marking point (with relevant number)
ecf	error carried forward
I	ignore
AVP	alternative valid point (examples given as guidance)

Page 3	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

1 (a) transport proteins – Y ;

pigments – X ;

(b) DNA

codes for, proteins / polypeptides / enzymes ;

one example of protein or enzyme ;

e.g. rubisco / electron acceptor / ATP synthase / transport

ref. transcription / mRNA ;

[max 2]

(c)

factor	stage	✓ or ✗
carbon dioxide concentration	Calvin cycle	✓
	photolysis	✗
light intensity	Calvin cycle	✗
	photolysis	✓
temperature	Calvin cycle	✓
	photolysis	✗

;;;

all 6 correct = 3 marks

4 or 5 correct = 2 marks

2 or 3 correct = 1 mark

[3]

[Total:7]

2 (a) presence of C base in DNA (code) changes amino acid (in myostatin) ;

myostatin in CC horses, is inactive / not produced ; **ora for TT**

in CC horses muscle, differentiation / growth, has not been slowed ; **ora for TT**

[max 2]

Page 4	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

(b) CC genotype does best in short races ;

data quote ; e.g. 75% winners at 1.0 km ;

TT genotype does better in longer races ;

data quote ; e.g. about 60% of winners at 2.6 km

CT genotype has some winners at all distances ;

CT does best at 1.8 km ;

[max 4]

(c) (i) by humans ;

individuals with desired features chosen to breed / AW / named example ;

[2]

(ii) can choose parents genotypes to breed ;

CC for racing short distances / TT for racing long distances / CT as all-rounders ;

ref. need to keep all three genotypes in population ;

[max 2]

[Total:10]

3 (a) (i) idea of sugars unable to pass through phospholipid bilayer ;

hydrophilic / polar / not lipid-soluble / water soluble ;

large ;

[max 2]

(ii) forms bonds with hydrophilic heads (of phospholipids) ;

hydrophobic parts of SWEET ;

bond with, fatty acid chains / hydrophobic tails, (of phospholipids) ;

ref. hydrogen bonding / ionic bonds / hydrophobic interactions ;

[max 3]

(b) (i) (SWEET) gene cannot be switched on ;

no SWEET (protein) produced ;

no, glucose / sugar, secreted (into intercellular spaces) ;

(so) Xoo / bacteria, do not multiply / numbers remain low ;

(small numbers of Xoo / bacteria) so no disease ;

[max 3]

Page 5	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

(ii) allele is recessive ;

idea of not expressed when dominant allele present ;

ref. promoter ; e.g. normal promoter must be inactivated or removed/must transfer mutated promoter

[max 2]

(iii) prevents diffusion of air (from leaves to roots) ;

ref. aerenchyma ;

roots respire anaerobically ;

(so) less ATP produced (for growth) ;

bacteria use of oxygen ;

more ethanol produced may be beyond tolerance/ **AW** ;

[max 4]

[Total:14]

4 (a) details of electrophoresis ;;

any 2 from

DNA cut by, restriction enzyme(s)/endonuclease(s)

loaded (into wells) at, negative end/cathode end, (of gel)

ref. buffer/electrolyte

(negatively charged) DNA attracted to, anode/positive electrode

separation due to, electric field/potential difference

short pieces/smaller mass, move further (in unit time)/move faster **ora**

fluorescent/radioactive, DNA probes

compare, DNA sequences/bands, (of male lizard and hatchling) ;

[3]

(b) (i) *body length*

no relationship (between body length and number of offspring) ;

small/intermediate, body length produce more offspring ;

sprint speed

lizards with greater sprint speed sire more offspring ;

use of two paired figures from Fig. 4.3 to support relationship ;

[4]

Page 6	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

- (ii) lizards with longer (hind) legs will tend to have more offspring ;
 so will have a greater chance of passing on their alleles ;
 (over time) the frequency of alleles (for long hind leg) will increase ;
 so mean hind leg length will increase ;
 directional selection ;

[max 3]

- (c) no, breeding / allele flow / gene flow, between (lizard) populations ;
 different selection pressures / different (environmental) conditions ;
 mutations occur ;
 advantageous alleles, selected for / passed on ;
 change in, allele frequency / gene pool ;
 genetic drift ;
 (eventually) unable to interbreed ;
allopatric speciation ;

[max 5]

[Total:15]

- 5 (a) anterior pituitary ;

[1]

- (b) (i) early follicle development not dependent on FSH ;
 with no FSH / no FSH receptors, follicle development stops ; **ora**
 with no FSH / no FSH receptors, Graafian / ovarian, follicle does not develop
 ; **ora**
 with no FSH / no FSH receptors, there is no ovulation ;
 no corpora lutea because these form, from Graafian follicle / after ovulation ;

[max 4]

Page 7	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

- (ii) sperm development better when FSH present ;

with FSH receptors

more sperm produced ; **ora**

sperm more active ; **ora**

males have increased fertility ; **ora**

without FSH receptors

some sperm produced ;

[max 3]

[Total:8]

- 6 (a) (i) tendency of water molecules to move from one region to another/potential energy of water/ability of water to do work ;

[1]

- (ii) (water potential) becomes, lower / more negative ;

[1]

- (iii) posterior pituitary ;

[1]

- (iv) for **one** mark ;

any 2 from

urine

sweat

water vapour (from exhaled air)

faeces

bleeding

tears

[max 1]

- (b) affects collecting duct, (cells/wall) ; **A** distal convoluted tubule cells

binds to receptors on cell surface membranes ;

activates series of enzyme controlled reactions ;

(phosphorylase causes), vesicles/aquaporins, to move to cell surface membrane (on lumen side) ;

vesicles/aquaporins, fuse with cell surface membrane ;

cells/wall, more permeable to water ;

water moves out of lumen (of collecting duct) ;

down water potential gradient ;

[max 5]

Page 8	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

- (c) produce, a lot of urine / dilute urine ;
 dehydration / thirsty ;
 cramps / loss of salts ;

[max 2]

[Total:11]

- 7 (a) symbols and key ; e.g. A = NF allele and a = normal allele

parental genotypes **and** gametes ; e.g. parental genotypes Aa x aa
 gametes A a x a a

offspring genotypes **and** phenotypes **linked** ; e.g. Aa has NF and aa is unaffected

[3]

- (b) spontaneous / random / chance ;

mutation of, gene / allele ;

AVP ; e.g. named mutagen / detail of mutation / in oocyte / in sperm

[max 2]

- (c) compresses nerve ;

damages, myelin sheaths / Schwann cells ;

prevents, setting up of local circuits / saltatory conduction ;

stops Na⁺ / K⁺ pumps from working ;

blocks blood supply;

qualified ; e.g. effect on, oxygen supply / glucose supply / ATP production

AVP ; e.g. may stop ion channels opening

[max 3]

[Total:8]

Page 9	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

8 (a) (i) receptors / hypothalamus, detect change in blood temperature ;

brain ;

(receptor / brain) sends impulses to effector ;

effector carries out response / example of response ;

blood temperature returns to normal ;

negative feedback ;

[max 4]

(ii) larger SA:V ratio ;

lose (relatively) more heat ;

ref. more mitochondria to release heat energy ;

cannot carry out behavioural actions to get warm ;

infants cannot shiver ;

[max 2]

(b) (i) **A** – ATP synthase / ATP synthetase / stalked particles ; **R** ATPase

B – inner membrane / crista ; **I** phospholipid bilayer

[2]

(ii) arrow going down from intermembrane space to matrix ;

[1]

(iii) **1 and 3** ;

[1]

(iv) water ;

[1]

(v) fatty acids ; **A** lipid / fat / triglycerides

[1]

[Total:12]

Page 10	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

- 9 (a) multicellular ;
- differentiated cells ;
- (most) have, vascular tissue / xylem and phloem ;
- eukaryotic (cells) ;
- ref. meristems ;
- (most) are not motile ;
- motile gametes **only** in mosses and ferns ;
- autotrophic nutrition / photosynthesis ;
- cells have:*
- chloroplasts ;
- large / central, vacuole ;
- walls made of cellulose ;

[max 8]

- (b) place in zoos ;
- protected against, disease / predation ;
- captive breeding programme ;
- ref. assisted reproduction / cloning / sperm banks ;
- released into wild ;
- ref. national parks / reserves ;
- rangers patrol parks ;
- human access restricted ;
- controlled agriculture ;
- controlled industry ;
- visitor centres / education ;
- habitat / breeding sites, protected ;
- banning sale of protected animals or their products ;
- banning hunting ;

[max 7]

[Total:15]

Page 11	Mark Scheme	Syllabus
	GCE AS/A LEVEL – May/June 2014	9700

10 (a) use Penicillium (in batch fermenter) ;

(main) nutrients added at start ;

penicillin only produced, after growth phase / when running out of nutrients ;

(penicillin) is a secondary metabolite ;

fermentation is stopped ;

penicillin is harvested ;

fermenter is cleaned out / ref. sterility ;

new culture of *Penicillium* is put in and started again ;

ref. fed batch culture ;

carbohydrate / named nutrient, added regularly ;

keeps fermentation going longer / produces more penicillin ;

detail of fermenter ;; e.g. paddle to mix nutrients / sterilising steam inlet
/ set at pH 6.5 / aeration / kept at 27 °C

[max 8]

(b) *batch*

easy to set up culture ;

can continue with minimal attention / AW ;

environmental conditions easy to control ;

fermenter can be used for different process afterwards ;

only waste one batch if contaminated ;

less chance of blockage in fermenter ;

continuous

no / less, down time / AW ;

small vessels can be used ;

productivity high ;

cost effective ;

downstream processing easier ;

good for using immobilised enzymes ;

[max 7]

[Total:15]