

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

* 0 5 5 7 6 0 8 7 6 *
* 0 5 5 7 6 0 8 7 6 *

CHEMISTRY

0620/33

Paper 3 (Extended)

October/November 2015

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **12** printed pages.

- 1 (a) Describe a chemical test which shows the presence of water.

test

colour change if water is present

..... [3]

- (b) How could you show that a sample of water is pure?

..... [1]

- (c) Describe how water is treated before it is supplied to homes and industry.

.....

..... [2]

- (d) State **two** industrial uses of water.

.....

..... [2]

[Total: 8]

- 2 Choose from the following list of gases. A gas may be chosen once, more than once or not at all.

sulfur dioxide

hydrogen

methane

carbon monoxide

argon

ethene

butane

- (a) It is used to bleach wood pulp. [1]

- (b) When burned in oxygen, the only product is water. [1]

- (c) It can polymerise. [1]

- (d) It is used to provide an inert atmosphere for welding. [1]

- (e) When reacted with oxygen, the only product is carbon dioxide. [1]

- (f) It is produced by the decay of vegetation in the absence of oxygen. [1]

[Total: 6]

3 Lithium bromide is an ionic compound. It can be electrolysed when it is molten or in aqueous solution. It cannot be electrolysed as a solid.

(a) Solid lithium bromide is a poor conductor of electricity. The ions cannot move to the electrodes, they are held in an ionic lattice by strong forces.

(i) Describe the motion of the ions in the solid state.

..... [1]

(ii) Define the term *ionic bonding*.

.....


..... [2]

(iii) What is meant by the term *ionic lattice*?

.....

..... [2]

(b) The diagram shows the electrolysis of molten lithium bromide.



(i) Mark on the diagram the direction of the electron flow. [1]

(ii) Write an ionic equation for the reaction at the negative electrode (cathode).

..... [1]

(iii) Write an ionic equation for the reaction at the positive electrode (anode).

..... [2]

(iv) Which ion is oxidised? Explain your answer.

.....

..... [2]

- (c) When aqueous lithium bromide is electrolysed, a colourless gas is formed at the negative electrode and the solution becomes alkaline.

Explain these observations and include an equation in your explanation.

.....

.....

.....

..... [3]

[Total: 14]

- 4 Two homologous series of hydrocarbons are the alkanes and the alkenes.

- (a) (i) One general characteristic of a homologous series is that the physical properties vary in a predictable way.

State **three** other general characteristics of a homologous series.

.....

.....

..... [3]

- (ii) How can the molecular formula of a hydrocarbon show whether it is an alkane or an alkene?

.....

..... [2]

- (iii) How do alkanes and alkenes differ in their molecular structures?

.....

..... [2]

(b) Cracking is the thermal decomposition of alkanes into smaller hydrocarbons and possibly hydrogen.

(i) State **two** conditions required for the cracking of an alkane.

..... [2]

(ii) One type of cracking produces an alkane and an alkene.

Complete an equation for the cracking of heptane into an alkane and an alkene.



(iii) Complete an equation for the cracking of heptane into hydrogen and two other products.



(iv) Suggest **one** reason why cracking is important.

..... [1]

(c) Hydrocarbons burn in excess oxygen to form carbon dioxide and water. 20 cm³ of a gaseous hydrocarbon burned in an excess of oxygen, 200 cm³. After cooling, the volume of the residual gas at r.t.p. was 150 cm³, 50 cm³ of which was oxygen.

(i) Determine the volume of the oxygen used.

..... [1]

(ii) Determine the volume of the carbon dioxide formed.

..... [1]

(iii) The hydrocarbon was an alkane.

Determine the formula of the hydrocarbon.

[1]

[Total: 15]

5 Sulfuric acid is a strong acid. In aqueous solution, it ionises as shown below.



(a) (i) What is meant by the term *acid*?

..... [1]

(ii) Sulfurous acid, H_2SO_3 , is a weak acid.

State the difference between a weak acid and a strong acid.

.....
 [2]

(b) Sulfurous acid forms salts called sulfites, which contain the ion SO_3^{2-} .

When barium nitrate solution is added to aqueous sulfurous acid, a white precipitate, **A**, forms.

Bromine water changes from brown to colourless when added to aqueous sulfurous acid.

Bromine oxidises sulfurous acid. When this solution is tested with acidified barium nitrate solution, a different white precipitate, **B**, is formed.

(i) Identify the white precipitate, **A**.

..... [1]

(ii) Identify the white precipitate, **B**.

..... [1]

(iii) Write an ionic equation for the reduction of the bromine molecule.

..... [1]

(iv) Name the product formed by the oxidation of sulfurous acid.

..... [1]

(c) Complete the following word equations.

(i) magnesium hydroxide + dilute sulfuric acid

..... [1]

(ii) zinc + dilute sulfuric acid

..... [1]

(iii) copper carbonate + dilute sulfuric acid

..... [1]

(d) Write equations for the reaction of dilute sulfuric acid with each of the following.

(i) ammonia

..... [2]

(ii) sodium hydroxide

..... [2]

(iii) iron

..... [2]

[Total: 16]

6 A reactivity series of metals is given below.

	metal name	symbol
most reactive ↓ least reactive	sodium	Na
	lithium	Li
	magnesium	Mg
	zinc	Zn
	manganese	Mn
	iron	Fe
	copper	Cu
	rhodium	Rh

(a) Which **two** metals will react most vigorously with cold water?

..... [1]

(b) Which **two** metals will not react with dilute hydrochloric acid?

..... [1]

(c) Deduce the formula of iron(III) sulfate.

..... [1]

(d) What is the formula of a magnesium ion?

..... [1]

(e) Describe a test-tube experiment which will show that manganese is more reactive than copper.

.....

 [3]

(f) Manganese is a typical transition metal.

Predict **three** physical and **two** chemical properties of this metal.

physical properties

.....
.....
.....


chemical properties

.....
.....


[5]

[Total: 12]

7 Two salts can be made from potassium hydroxide and sulfuric acid. They are potassium sulfate, K_2SO_4 , and the acid salt potassium hydrogen sulfate, $KHSO_4$. They are both made by titration.



(a) 25.0 cm³ of potassium hydroxide, concentration 2.53 mol/dm³, was neutralised by 28.2 cm³ of dilute sulfuric acid.



Calculate the concentration of the sulfuric acid.

number of moles of KOH used =

number of moles of H₂SO₄ needed to neutralise the KOH =

concentration of dilute sulfuric acid = mol/dm³


[3]

(b) In the conical flask there is a neutral solution of potassium sulfate which still contains the indicator used in the titration.

(i) Describe how you could obtain a solution of potassium sulfate without the indicator.

.....
 [2]

(ii) Potassium hydrogen sulfate can be made by the following reaction.



Suggest how you could make a solution of potassium hydrogen sulfate without using an indicator.

.....

 [2]

(c) Describe a test which would distinguish between aqueous solutions of potassium sulfate and sulfuric acid.

test

result

[2]

[Total: 9]

DATA SHEET
The Periodic Table of the Elements

		Group																										
		I	II	III	IV	V	VI	VII	VIII	IX	X																	
		<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">1 H Hydrogen 1</div> <div style="border: 1px solid black; padding: 2px;">4 He Helium 2</div> </div>																										
3	7	9	23	24	39	40	85	133	226	87	88	89																
	Li Lithium 4	Be Beryllium 9	Na Sodium 11	Mg Magnesium 12	K Potassium 19	Ca Calcium 20	Rb Rubidium 37	Cs Caesium 55	Fr Francium 87	Ba Barium 56	Sr Strontium 38	Y Yttrium 39	La Lanthanum 57	Ra Radium 88	Ac Actinium 89													
		48	51	52	55	56	59	59	64	65	70	73	75	79	80	84												
		Ti Titanium 22	V Vanadium 23	Cr Chromium 24	Mn Manganese 25	Fe Iron 26	Co Cobalt 27	Ni Nickel 28	Cu Copper 29	Zn Zinc 30	Ga Gallium 31	Ge Germanium 32	As Arsenic 33	Se Selenium 34	Br Bromine 35	Kr Krypton 36												
		91	93	96	101	103	106	108	112	115	119	122	127	128	131	137												
		Zr Zirconium 40	Nb Niobium 41	Mo Molybdenum 42	Tc Technetium 43	Ru Ruthenium 44	Rh Rhodium 45	Pd Palladium 46	Ag Silver 47	Cd Cadmium 48	In Indium 49	Sn Tin 50	Sb Antimony 51	Te Tellurium 52	I Iodine 53	Xe Xenon 54												
		178	181	184	186	190	192	195	197	201	204	207	209	210	210	210												
		Hf Hafnium 72	Ta Tantalum 73	W Tungsten 74	Re Rhenium 75	Os Osmium 76	Ir Iridium 77	Pt Platinum 78	Au Gold 79	Hg Mercury 80	Tl Thallium 81	Pb Lead 82	Bi Bismuth 83	Po Polonium 84	At Astatine 85	Rn Radon 86												
		140	141	144	150	152	157	159	162	165	167	169	173	175	232	238												
		Ce Cerium 58	Pr Praseodymium 59	Nd Neodymium 60	Sm Samarium 62	Pm Promethium 61	Gd Gadolinium 64	Tb Terbium 65	Dy Dysprosium 66	Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71	Th Thorium 90	U Uranium 92	Pa Protactinium 91	Np Neptunium 93	Pu Plutonium 94	Am Americium 95	Cm Curium 96	Bk Berkelium 97	Cf Californium 98	Es Einsteinium 99	Fm Fermium 100	Md Mendelevium 101	No Nobelium 102	Lr Lawrencium 103

*58-71 Lanthanoid series
†90-103 Actinoid series

Key

a	X
b	†

a = relative atomic mass
X = atomic symbol
b = proton (atomic) number

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).