



Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			0620/41
Paper 4 Theory	/ (Extended)	Octol	oer/November 2017
			1 hour 15 minutes
Candidates ans	swer on the Question Paper.		

READ THESE INSTRUCTIONS FIRST

No Additional Materials are required.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



1 The table gives information about five particles. The particles are all atoms or ions.


particle	number of protons	number of neutrons	number of electrons
Α	6	8	6
В	12	12	12
С	13	14	10
D	8	8	10
E	11	12	11

Answer the following questions using the information in the table. Each particle may be used once, more than once or not at all.

(a) Which particle, A, B, C, D or E,

	(i)	is an atom with atomic number 12,	
			[1]
	(ii)	is an atom with nucleon number 14,	
			[1]
	(iii)	is an ion with a positive charge,	
	·· \		[1]
	(iv)	has only one electron in its outer shell?	[4]
			ניו
(b)	D is	s an ion of an element.	
	lder	ntify the element and write the formula of D .	
			[2]
		[Total:	6]

2 The graph shows how the temperature of a substance changes as it is cooled over a period of 30 minutes. The substance is a gas at the start.



Each letter on the graph may be used once, more than once or not at all.

(a)	Which letter	. S.	Τ.	. V.	W.	X.	Υ	or Z	. shows	when

	(i)	the particles in the substance have the most kinetic energy,
		[1
	(ii)	the particles in the substance are furthest apart,
		[1
((iii)	the substance exists as both a gas and a liquid?
		[1
(b)	Use	the graph to estimate the freezing point of the substance.
		°C [1
(c)	Nar	ne the change of state directly from a solid to a gas.
		[1
(d)	Who	en smoke is viewed through a microscope, the smoke particles in the air appear to jumpend.
	(i)	What term describes this movement of the smoke particles?
		[1
	(ii)	Explain why the smoke particles move in this way.

3	(a)	When magnesium is added to aqueous copper(II) sulfate a reaction occurs.
		The ionic equation for the reaction is shown.


$Mg + Cu^{2+} \rightarrow Mg^{2+} + Cu$	Ma	+	Cu ²⁺	\rightarrow	Ma ²⁺	+	Cu
---	----	---	------------------	---------------	------------------	---	----

((i)	Give one change you would observe during this reaction.	[4]
(i	ii)	Explain why this is a redox reaction.	ניו
(ii	ii)	Identify the oxidising agent in this reaction. Give a reason for your answer.	
			[2]
(i	v)	A redox reaction occurs when magnesium is heated with iron(III) oxide.	
		Write a chemical equation for the reaction between magnesium and iron(III) oxide.	
		metal iron and the alloy steel are commonly used materials. A problem with them is t	
	(i)	How does painting iron and steel prevent rusting?	
			[1]
(i	ii)	Magnesium blocks can be attached to the bottom of steel boats.	
		Explain how the magnesium blocks prevent the whole of the bottom of the boat frusting.	om


(iii)	Replacing the magnesium blocks with copper blocks does not prevent rusting.
	Explain why the copper blocks do not prevent rusting.
	[1]
	[Total: 10]

(a) Eth	nanol, C ₂ H ₅ OH, can be made by fermentation.	
(i)	Complete the chemical equation for the formation of ethanol by fermentation.	
	$C_6H_{12}O_6 \rightarrowC_2H_5OH +$	[2]
(ii)	State two conditions required for fermentation.	[-]
	1	
	2	 [2]
	nanol can also be made by the catalytic hydration of ethene. The equation for the reaction own.	ı is
	$C_2H_4 + H_2O \rightarrow C_2H_5OH$	
(i)	Name a suitable catalyst for this reaction.	F41
(ii)	Calculate the maximum mass of ethanol that can be made from 56 g of ethene.	[1]
	maximum mass of ethanol = g	וכו
(-) - 4		[4]
	nanol can be oxidised to form ethanoic acid.	
(i)	Name a suitable oxidising agent for this reaction.	
		[1]

(ii) A molecule of ethanoic acid has the structure shown.



Complete the dot-and-cross diagram to show the electron arrangement in ethanoic acid. Show outer shell electrons only.



[3]

(d) Ethanoic acid is a weak acid.

(i)	When referring to an acid, what is meant by the term weak?
	[1]
(ii)	Describe how you could show that ethanoic acid is a weaker acid than hydrochloric acid.
	[3]

(e) Carboxylic acids react with alcohols to make esters.

The structure of an ester is shown.

Draw the structures of the carboxylic acid and alcohol from which this ester can be made. Give the names of the carboxylic acid and alcohol.

structure of the carboxylic acid

name of the carboxylic acid	

structure of the alcohol

name of the alcohol[4]

[Total: 19]

5

(a)		id copper(II) carbonate undergoes thermal decomposition. One of the products of rmal decomposition is copper(II) oxide.	the
	(i)	State the colour change of the solid seen during the reaction.	
		start colour	
		end colour	
	(ii)	Write a chemical equation for the thermal decomposition of copper(II) carbonate.	[1]
			[1]
(b)		$\mbox{\rm oper}(II)$ carbonate reacts with dilute nitric acid. One of the products of the reaction ution of $\mbox{\rm copper}(II)$ nitrate.	is a
	(i)	Describe tests for $copper(II)$ ions and nitrate ions. Include the results of the tests.	
		copper(II) ions	
		nitrate ions	
			[4]
	(ii)	Copper(II) nitrate undergoes thermal decomposition.	
		Balance the chemical equation for the thermal decomposition of $copper(II)$ nitrate.	
		Cu(NO ₃) ₂ \rightarrow CuO +NO ₂ +O ₂	[1]


(c) Nitrogen dioxide, NO_2 , exists in equilibrium with dinitrogen tetroxide, N_2O_4 . Nitrogen dioxide is brown and dinitrogen tetroxide is colourless.

$$2NO_2(g) \rightleftharpoons N_2O_4(g)$$

brown colourless

(i) A sample of nitrogen dioxide and dinitrogen tetroxide at equilibrium was placed in a closed gas syringe.

The syringe plunger was pushed in. This increased the pressure in the gas syringe. The temperature was kept constant.




	the position of the equilibrium.
	[3]
(ii)	A sealed tube containing nitrogen dioxide and dinitrogen tetroxide at equilibrium was cooled in an ice bath at constant pressure. The contents of the tube became paler.
	Suggest an explanation for this observation in terms of the position of the equilibrium.
	[2]

[Total: 12]

6

(ii) The melting point of aluminium oxide is above 2000 °C.		11	
(i) Complete the dot-and-cross diagram to show the electron arrangement in one of the oxide ions present in aluminium oxide. Include the charge on the oxide ion. One of the aluminium ions is shown. 3+	Alumin	ium is extracted from aluminium oxide by electrolysis.	
(b) Aluminium oxide is an ionic compound with a high melting point. (i) Complete the dot-and-cross diagram to show the electron arrangement in one of the oxide ions present in aluminium oxide. Include the charge on the oxide ion. One of the aluminium ions is shown. 3+	(a) Wh	hy is aluminium not extracted by heating aluminium oxide with carbon?	
(i) Complete the dot-and-cross diagram to show the electron arrangement in one of the oxide ions present in aluminium oxide. Include the charge on the oxide ion. One of the aluminium ions is shown. 3+			 [1]
oxide ions present in aluminium oxide. Include the charge on the oxide ion. One of the aluminium ions is shown.	(b) Alu	uminium oxide is an ionic compound with a high melting point.	
(ii) The melting point of aluminium oxide is above 2000 °C.	(i)	oxide ions present in aluminium oxide. Include the charge on the oxide ion.	the
(ii) The melting point of aluminium oxide is above 2000 °C.			
			[2]
	(ii)	The melting point of aluminium oxide is above 2000 °C.	
Explain why aluminium oxide has a high melting point.		Explain why aluminium oxide has a high melting point.	

(c) Aluminium can be extracted by electrolysis using the apparatus shown.



	(i)	Name the type of particle responsible for the transfer of charge in	
		the wires,	
		the electrolyte.	 2]
	(ii)	Give two reasons why cryolite is used. 1	-
		2	
			2]
	(iii)	Write the ionic half-equation for the formation of aluminium during the electrolysis.	
		[1]
	(iv)	Explain how carbon dioxide gas is formed at the anodes.	
		[3	3]
(d)	rea If th	en a piece of aluminium is placed in dilute hydrochloric acid, there is no immediate visiblection. The aluminium is left in the dilute hydrochloric acid for several hours, bubbles start to form. Totalin why aluminium does not react immediately with dilute hydrochloric acid.	
		[´	1]

[Total: 14]

Question 7 starts on the next page.

7 Copper(II) oxide reacts with dilute hydrochloric acid.

$$CuO(s) + 2HCl(aq) \rightarrow CuCl_2(aq) + H_2O(l)$$

 $6.00\,g$ of copper(II) oxide were added to $50.0\,cm^3$ of $1.00\,mol/dm^3$ hydrochloric acid. This was an excess of copper(II) oxide.

(a)	The rate of the reaction can be increased by increasing the concentration of the hydrochloric acid
	or by heating it.

(i)	In terms of collisions, explain why increasing the concentration of the hydrochloric acid increases the rate of the reaction.
	[2]
(ii)	In terms of collisions, explain why heating the hydrochloric acid increases the rate of the reaction.
	[0]

(b) (i) Calculate the number of moles of copper(II) oxide added to the hydrochloric acid.	
moles of copper(II) oxide = (ii) Calculate the number of moles of hydrochloric acid used.	mol [2]
moles of hydrochloric acid = (iii) Calculate the mass of copper(II) oxide that did not react.	mol [1]
mass of copper(II) oxide that did not react =	g [2]
(c) Crystals of hydrated copper(${\rm II}$) chloride were obtained from the solution at the end reaction.	of the
The crystals had the following composition by mass: Cl, 41.52%; Cu, 37.43%; H, O, 18.71%.	2.34%;
Calculate the empirical formula of the crystals.	
empirical formula =	[2]
TC]	otal: 11]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

	III	2 He	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	R	radon			
				6	щ	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	Н	iodine 127	85	Ą	astatine -			
	5			80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	Тe	tellurium 128	84	Po	molonium —	116	_	livermorium —
	>			7	z	nitrogen 14	15	₾	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	Ξ	bismuth 209			
	≥			9	ပ	carbon 12	14	S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	Ρl	flerovium -
	≡			2	Δ	boron 11	13	Αſ	aluminium 27	31	Ga	gallium 70	49	In	indium 115	84	lT	thallium 204			
										30	Zn	zinc 65	48	g	cadmium 112	80	Я	mercury 201	112	ပ်	copernicium -
										59	on	copper 64	47	Ag	silver 108	62	Au	gold 197	111	Rg	roentgenium -
Group										28	Z	nickel 59	46	Pd	palladium 106	78	₽	platinum 195	110	Ds	darmstadtium -
يَّق				,						27	ဝိ	cobalt 59	45	格	rhodium 103	77	'n	iridium 192	109	Ĭ	meitnerium -
		- エ	hydrogen 1							26	Fe	iron 56	44	Ru	ruthenium 101	92	Os	osmium 190	108	Hs	hassium –
							1			25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium —
				_	pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	>	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	q	niobium 93	73	Б	tantalum 181	105	o O	dubnium —
					atc	rel				22	j	titanium 48	40	Zr	zirconium 91	72	茔	hafnium 178	104	Ŗ	rutherfordium -
										21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ba	barium 137	88	Ra	radium –
	_			က	=	lithium 7	11	Na	sodium 23	19	×	potassium 39	37	ВВ	rubidium 85	22	Cs	caesium 133	87	ь Г	francium -

77	lutetium 175	103	ב	lawrencium	ı
0 5	ytterbium 173	102	%	nobelium	ı
69 L	thulium 169	101	Md	mendelevium	I
89 7	erbium 167	100	Fm	ferminm	1
⁶⁷	holmium 165	66	Es	einsteinium	I
99	dysprosium 163	86	ŭ	californium	ı
65 Th	terbium 159	97	BK	berkelium	1
49 C	gadolinium 157	96	Cm	curium	ı
63 <u>T</u>	europium 152	92	Am	americium	ı
62 An	samarium 150	94	Pu	plutonium	ı
61 D	promethium	93	δ	neptunium	ı
09 Z	neodymium 144	92	\supset	uranium	238
59 7	praseodymium 141	91	Ра	protactinium	231
28 0	cerium 140	06	Ч	thorium	232
57	lanthanum 139	68	Ac	actinium	I

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).