

9

	MANN. Daha
	UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education
CANDIDATE NAME	
CENTER NUMBER	CANDIDATE NUMBER
MATHEMATIC	S (US) 0444/43
Paper 4 (Exten	ded) October/November 2013
	2 hours 30 minutes
Candidates ans	wer on the Question Paper.

Additional Materials: Geometrical instruments Electronic calculator

READ THESE INSTRUCTIONS FIRST

Write your Center number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

If work is needed for any question it must be shown in the space provided.

Electronic calculators should be used.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant digits.

Give answers in degrees to one decimal place.

For π , use either your calculator value or 3.142.

The number of points is given in parentheses [] at the end of each question or part question. The total of the points for this paper is 130.


Write your calculator model in the box below.

This document consists of **19** printed pages and **1** blank page.




Formula List

	2	man.	Dapa Cambridge.com
	Formula L	ist	PaCan .
For the equation	$ax^2 + bx + c = 0$	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$	"Bridge.c
Lateral surface area, A, of cylin	nder of radius r, height h.	$A = 2\pi rh$	9177
Lateral surface area, A, of cond	e of radius r, sloping edge l.	$A = \pi r l$	
Surface area, A, of sphere of ra	adius <i>r</i> .	$A = 4\pi r^2$	
Volume, <i>V</i> , of pyramid, base an	rea A , height h .	$V = \frac{1}{3}Ah$	
Volume, V , of cone of radius r ,	, height <i>h</i> .	$V = \frac{1}{3}\pi r^2 h$	
Volume, <i>V</i> , of sphere of radius	r.	$V = \frac{4}{3}\pi r^3$	
		$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ $a^2 = b^2 + c^2 - 2bc \cos A$	



 $2bc \cos A$ $= b^2 + c^2$ Area = $\frac{1}{2}bc\sin A$




www.papacambridge.com 4 2 A field, *ABCD*, is in the shape of a quadrilateral. A footpath crosses the field from *A* to *C*. C26 В NOT TO SCALE 55 m 65° . 122° 32° D 62 m (a) Use the sine rule to calculate the distance AC and show that it rounds to 119.9 m, correct to 1 decimal place.


Answer(a)

(b) Calculate the length of *BC*.


Answer(b) m [4]


[3]






(d) On the grid opposite, draw the graph of $V = 4x^3 - 32x^2 + 63x$ for $0 \le x \le 3.5$. Three of the points have been plotted for you.











	10 ney spent on t	he internet in	one day.	ANNA DODO	For miner's
$1 1 < t \le 2$	$2 < t \le 3$	$3 < t \le 5$	$5 < t \le 7$	$7 < t \le 10$	ennihridge con
11	10	19	13	12	COM
1	$1 1 < t \le 2$	Its. $1 1 < t \le 2 2 < t \le 3$	Its. 1 $1 < t \le 2$ $2 < t \le 3$ $3 < t \le 5$	1 $1 < t \le 2$ $2 < t \le 3$ $3 < t \le 5$ $5 < t \le 7$	1 $1 < t \le 2$ $2 < t \le 3$ $3 < t \le 5$ $5 < t \le 7$ $7 < t \le 10$













The cross section of the container is designed from three semi-circles with diameters 17.5 cm, 6.5 cm and 24 cm.




(a) Calculate the area of the cross section of the container.


(b) Calculate the external surface area of the container, including the base.


Answer(b) cm² [4]





 $Answer(d)(ii) h = \dots cm [3]$












The diagrams are made from dots (\bullet) and squares (\Box) joined by lines.

(a) Complete the table.

Diagram	1	2	3	4	5	п	
Number of dots	6	9	12				
Number of squares	0	1	3			$\frac{1}{2}n(n-1)$	
Number of triangles	4	9	16				
Number of lines	9	18	30	45	63	$\frac{3}{2}(n+1)(n+2)$	

(b) Which diagram has 360 lines?



- (c) The total number of lines in the first *n* diagrams is
 - (i) When n = 1, show that $p + q = 8\frac{1}{2}$. Answer(c)(i)
 - (ii) By choosing another value of n and using the equation in part (c)(i), find the values of p and q.

 $Answer(c)(ii) p = \dots$

19

 $\frac{1}{2}n^3 + pn^2 + qn.$



BLANK PAGE

20

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.