



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

| CANDIDATE NAME | | | | |
|-------------------|--|---------------------|--|--|
| CENTRE NUMBER | | CANDIDATE NUMBER | | |

PHYSICAL SCIENCE

0652/51

Paper 5 Practical Test

October/November 2012

1 hour 30 minutes

Candidates answer on the Question Paper.

Additional Materials:

As listed in Instructions to Supervisors

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Chemistry practical notes for this paper are printed on page 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

| For Examiner's Use | |
|--------------------|--|
| 1 | |
| 2 | |
| Total | |

This document consists of 10 printed pages and 2 blank pages.



BLANK PAGE

1 (a) You are going to find the mass of a metre rule. Refer to Fig. 1.1 as you follow these instructions.

For Examiner's Use




Fig. 1.1

- (i) Clamp a knife edge or triangular piece of wood in a clamp on a stand; this will act as a pivot on which to balance the rule.
 - Place the metre rule horizontally on the pivot at the 40 cm mark on the rule.
 - Hang the 100g mass provided from the 10cm mark on the rule, using half of the thread.
 - Now hang a second mass, *m*, of 50 g, using the other half of the thread, from the rule on the other side of the pivot from the 100 g mass.
 - Adjust the distance, *d*, from the pivot to the 50 g mass by moving the 50 g mass to make the rule balance.
 - Measure the distance, *d*, from the pivot to the 50 g mass.

Record this value in Table 1.1 on page 4.

[1]

For Examiner's Use

| mass, <i>m</i> /g | distance, <i>dl</i> cm | $\frac{1}{\text{mass}, m} / \frac{1}{\text{g}}$ | mass, $m \times$ distance, d/g cm |
|----------------------|---------------------------|---|-------------------------------------|
| 50 | | 0.020 | |
| 60 | | | |
| 70 | | | |
| 80 | | | |
| 90 | | 0.011 | |

(ii) Repeat (i) using a 60 g mass instead of the 50 g mass to balance the rule, ensuring that the pivot is still at the 40 cm mark and the 100 g mass is still at the 10 cm mark.

Record the new value of distance, *d*, in Table 1.1.

(iii) Repeat (i) using 70 g, 80 g and 90 g masses to balance the rule, ensuring that the pivot is still at the 40 cm mark and the 100 g mass is still at the 10 cm mark.


Record the values of distance, *d*, in Table 1.1. [3]

(b) (i) Calculate $\frac{1}{\text{mass}, m}$ for the three remaining values of mass, m, and record your answers in Table 1.1. [1]

(ii) Complete the vertical scale and plot a graph of distance, d, against $\frac{1}{\text{mass}, m}$ on the grid provided.

For Examiner's Use

Draw the best straight line.



| | (iii) | Find the gradient of the line of your graph, showing clearly your working. | For Examiner's Use |
|-----|-------|--|--------------------------|
| | (iv) | $gradient = \underline{\hspace{1cm}} [2]$ Use the value of the gradient from (b)(iii) and the formula shown to calculate the mass of the rule. $mass of rule = 300 - (0.1 \times gradient)$ | |
| (c) | | mass of rule = $_{_{_{_{_{_{_{_{_{_{_{_{_{_{_{_{_{_{_{$ |] |
| | (ii) | [1] Calculate the average of your mass, $m \times$ distance, d values in Table 1.1. |] |
| | | average (mass, $m \times$ distance, d) = g cm [1 |] |

For Examiner's Use

| 1 | (iii) Use the value of the average from (c)(ii) and the formula to calculate the mass of the rule. |
|-----|--|
| | mass of rule = 300 - $\left(\frac{\text{average (mass, } m \times \text{distance, } d)}{10}\right)$ |
| | |
| | |
| | mass of rule = g [1] |
| (d) | What is the advantage of plotting a graph and finding the gradient rather than using the average (mass, m , × distance, d) value to calculate the mass of the rule? |
| | [1] |

2 You are going to investigate the acidity of a soil sample. The sample of soil has been shaken with purified water and filtered. You have been given the filtrate as soil washings, **S**.

For Examiner's Use

You are going to find the concentration of acid in **S**.

- (a) (i) Use a measuring cylinder to place 10 cm³ of the alkali aqueous calcium hydroxide into a small beaker.
 - Add 3 drops of indicator **P** to the alkali in the beaker.
 - Place 10 cm³ soil washings **S** in the rinsed 0-10 cm³ measuring cylinder.
 - Slowly add, drop by drop, soil washings S from the measuring cylinder to the beaker containing the alkali and the indicator P until the mixture just becomes colourless.

Record, in Table 2.1, the volume of unused soil washings, V_R , remaining in the measuring cylinder. [1]

Table 2.1

| volume of aqueous calcium hydroxide, V_a/cm^3 | volume of unused soil washings, V_R/cm^3 | volume of soil washings used, <i>V_s</i> /cm ³ |
|---|---|---|
| 10.0 | | |
| 10.0 | | |
| 10.0 | | |

- (ii) Repeat (a)(i) twice more, recording the volume of unused soil washings, V_R , remaining in the measuring cylinder in Table 2.1. [3]
- (iii) Calculate the volume of soil washings used, V_s , in each case and record the values in column three of Table 2.1. [1]
- (iv) Calculate the average volume of soil washings used, $V_{\rm av}$, using the data in Table 2.1.

 $V_{av} =$ cm³ [1]

For Examiner's Use

| (V) | | e the concentration, c_s , of the acid (H $$ ion) in the soil washings, $oldsymbol{s}$, usir wing formula and data | ıg |
|---------|----------|--|-----|
| | | $c_s \times V_{av} = 2 \times c_a \times V_a$ | |
| | | erage value from (a)(iv) centration of alkali, 0.013 mol/dm ³ cm ³ | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | $c_{\rm s}$ = mol/dm ³ [| [3] |
| (b) (i) | Place 5 | cm ³ soil washings, S , in a test-tube and add four drops of Univers | al |
| | Record t | the colour and pH of the soil washings. Keep the mixture for (b)(ii) . | |
| | colour = | | |
| | pH = | [| [1] |
| (ii) | | patula load of solid calcium carbonate to the mixture from (b)(i). Stopp tube and with your finger on the stopper shake well. | er |
| | Record t | the colour and pH of the resulting mixture. Keep this mixture for (b)(iii) . | |
| | colour = | | |
| | pH = | [| [1] |
| (iii) | | nixture from (b)(ii) add a second spatula load of solid calcium carbonat and shake. | e, |
| | Once ag | ain record the colour and pH of the resulting mixture. | |
| | colour = | | |
| | pH = | [| [1] |

(c) It is essential that you wear eye protection.

For Examiner's Use

- Place 5 cm3 soil washings, S, in a test-tube and add four drops of Universal Indicator.
- Add a spatula load of solid calcium hydroxide to the test-tube.
- Put a stopper in the test-tube and with your finger on the stopper, shake well.

| | Record tl | he colour and pH of the resulting mixture. | |
|-----|--|--|-----|
| | colour = | | |
| | pH = | | [1] |
| (d) | | which substance, calcium carbonate or calcium hydroxide, would require amount to spread on the soil to neutralise acidity in the soil. | the |
| | Explain your answer using the results from (b) and (c) . | | |
| | | | |
| | | | |
| | | | [2] |

BLANK PAGE

CHEMISTRY PRACTICAL NOTES

Test for anions

| anion | test | test result |
|---|---|--|
| carbonate (CO ₃ ²⁻) | add dilute acid | effervescence, carbon dioxide produced |
| chloride (Cl ⁻) [in solution] | acidify with dilute nitric acid, then add aqueous silver nitrate | white ppt. |
| nitrate (NO ₃ ⁻) [in solution] | add aqueous sodium hydroxide then aluminium foil; warm carefully | ammonia produced |
| sulfate (SO ₄ ²⁻) [in solution] | acidify then add aqueous barium chloride <i>or</i> aqueous barium nitrate | white ppt. |

Test for aqueous cations

| cation | effect of aqueous sodium hydroxide | effect of aqueous ammonia |
|--|--|--|
| ammonium (NH ₄ ⁺) | ammonia produced on warming | - |
| copper(II) (Cu ²⁺) | light blue ppt., insoluble in excess | light blue ppt., soluble in excess giving a dark blue solution |
| iron(II) (Fe ²⁺) | green ppt., insoluble in excess | green ppt., insoluble in excess |
| iron(III) (Fe ³⁺) | red-brown ppt., insoluble in excess | red-brown ppt., insoluble in excess |
| zinc (Zn ²⁺) | white ppt., soluble in excess giving a colourless solution | white ppt., soluble in excess giving a colourless solution |

Test for gases

| gas | test and test results | |
|-----------------------------------|----------------------------------|--|
| ammonia (NH ₃) | turns damp red litmus paper blue | |
| carbon dioxide (CO ₂) | turns limewater milky | |
| chlorine (Cl ₂) | bleaches damp litmus paper | |
| hydrogen (H ₂) | "pops" with a lighted splint | |
| oxygen (O ₂) | relights a glowing splint | |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.