



General Certificate of Secondary Education
2018

Statistics

Unit 1

Foundation Tier

[GST11]

THURSDAY 21 JUNE, MORNING

**MARK
SCHEME**

General Marking Instructions

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right hand column and they are prefixed by the letters **M**, **A** and **MA** as appropriate. The key to the mark scheme is given below:

- M** indicates marks for correct method.
- A** indicates marks for accurate working, whether in calculation, readings from tables, graphs or answers.
- MA** indicates marks for combined method and accurate working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be **followed through** from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

It should be noted that where an error trivialises a question, or changes the nature of the skills being tested, then as a general rule, it would be the case that not more than half the marks for that question or part of that question would be awarded; in some cases the error may be such that no marks would be awarded.

Positive marking

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examiner)

		AVAILABLE MARKS																
1	(a) 7	A1																
	(b) Dungannon: ♦ ♦ ♦	MA1																
	(c) Disagree Any valid reason	A1																
	• Raindrop symbol is symmetrical so can be easily adjusted to represent one day	A1																
		4																
2	(a) $80 - (12 + 19 + 6 + 13 + 14 + 9)$ $= 80 - 73$ $= 7$	M1 A1																
	(b) Tally	MA1																
	Frequency 7	A1																
	(c) Silver	A1																
	(d) Black	A1																
	(e)																	
	<table border="1"> <thead> <tr> <th>Colour of car</th> <th>Frequency</th> </tr> </thead> <tbody> <tr> <td>Black</td> <td>12</td> </tr> <tr> <td>Silver</td> <td>19</td> </tr> <tr> <td>Blue</td> <td>6</td> </tr> <tr> <td>Red</td> <td>13</td> </tr> <tr> <td>Green</td> <td>7</td> </tr> <tr> <td>White</td> <td>14</td> </tr> <tr> <td>Other</td> <td>9</td> </tr> </tbody> </table>	Colour of car	Frequency	Black	12	Silver	19	Blue	6	Red	13	Green	7	White	14	Other	9	MA1
Colour of car	Frequency																	
Black	12																	
Silver	19																	
Blue	6																	
Red	13																	
Green	7																	
White	14																	
Other	9																	
	(f) Primary data	A1																
		8																

					AVAILABLE MARKS
3	(a)				
		Biology	Chemistry	Physics	TOTAL
	Boys	35	39	48	122
	Girls	43	23	52	118
	TOTAL	78	62	100	240
				MA3	
	(b) $P(\text{Bio}) = \frac{78}{240} = \frac{13}{40} = 0.325$			MA1	
	(c) $P(\text{Not Chem} \mid \text{Girl}) = \frac{95}{118} = 0.805$			MA1	A1
	(d) Expected No. of Pupils = $24 \times \frac{100}{240}$ = 10			MA1	8
4	(a) The residents of the town			A1	
	(b) ‘Recently’ not specific enough/open to interpretation as no time frame No box for not visiting the library at all.			A1	
	(c) Any valid advantage, e.g. • Easy to analyse			A1	4
5	(a) 0	3	5		
	1	1	2	4	6
	2	1	5	7	8
	3	0	2	3	
	4	1			MA3
	Key: 0 3 represents 3 text messages			A1	
	(b) (i) Median = $\frac{1}{2}(17 + 21)$ = 19 text messages			M1	
	(ii) Range = $41 - 3$ = 38 text messages			MA1	
	(c) Any valid comparison involving the medians • Boys receive more messages than girls • Girls receive fewer texts than boys			A1	
	Any valid comparison involving the ranges • Number of texts is more variable/less consistent among girls			A1	9

			AVAILABLE MARKS
6	(a) (i) £5.80 (ii) $\text{£}7.30 - \text{£}4.00$ = £3.30	A1 M1 A1	
	(b) Values in correct places (ft their values) Correct shape (must have values marked)	MA1 MA1	
	(c) Negative skew Median closer to UQ (or equivalent)	A1 A1	
	(d) Method which uses people available at a particular place/time.	A1	8
7	(a) 4 years (b) Negative correlation (c) £5800 (d) (i) £11 000 to £11 200 (ii) May not be reliable as straight line model may not apply in first year.	A1 A1 MA1 MA1 A1	5
8	(a) (i) B (ii) C (iii) A (b) Coin is possibly biased as the number of ‘heads’ exceeds the expected number. (c) Repeat the experiment a larger number of times.	A1 A1 A1 A2 A1	6

				AVAILABLE MARKS																																	
9	(a)	Any advantage, e.g. • no need to have a column for every possible time	A1																																		
		Any disadvantage, e.g. • accuracy is lost by grouping	A1																																		
(b)		<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: center;">Time</th><th style="text-align: center;">Frequency, f</th><th style="text-align: center;">t</th><th style="text-align: center;">ft</th></tr> </thead> <tbody> <tr> <td style="text-align: center;">$0 < t \leq 10$</td><td style="text-align: center;">343</td><td style="text-align: center;">5</td><td style="text-align: center;">1715</td></tr> <tr> <td style="text-align: center;">$10 < t \leq 20$</td><td style="text-align: center;">214</td><td style="text-align: center;">15</td><td style="text-align: center;">3210</td></tr> <tr> <td style="text-align: center;">$20 < t \leq 30$</td><td style="text-align: center;">197</td><td style="text-align: center;">25</td><td style="text-align: center;">4925</td></tr> <tr> <td style="text-align: center;">$30 < t \leq 40$</td><td style="text-align: center;">145</td><td style="text-align: center;">35</td><td style="text-align: center;">5075</td></tr> <tr> <td style="text-align: center;">$40 < t \leq 50$</td><td style="text-align: center;">72</td><td style="text-align: center;">45</td><td style="text-align: center;">3240</td></tr> <tr> <td style="text-align: center;">$50 < t \leq 60$</td><td style="text-align: center;">29</td><td style="text-align: center;">55</td><td style="text-align: center;">1595</td></tr> <tr> <td style="text-align: center;">TOTAL</td><td style="text-align: center;">1000</td><td></td><td style="text-align: center;">19760</td><td></td></tr> </tbody> </table>	Time	Frequency, f	t	ft	$0 < t \leq 10$	343	5	1715	$10 < t \leq 20$	214	15	3210	$20 < t \leq 30$	197	25	4925	$30 < t \leq 40$	145	35	5075	$40 < t \leq 50$	72	45	3240	$50 < t \leq 60$	29	55	1595	TOTAL	1000		19760		MA2	
Time	Frequency, f	t	ft																																		
$0 < t \leq 10$	343	5	1715																																		
$10 < t \leq 20$	214	15	3210																																		
$20 < t \leq 30$	197	25	4925																																		
$30 < t \leq 40$	145	35	5075																																		
$40 < t \leq 50$	72	45	3240																																		
$50 < t \leq 60$	29	55	1595																																		
TOTAL	1000		19760																																		
		$\bar{t} = \frac{19760}{1000}$	M1																																		
		= 19.76 seconds	A1																																		
10	(c)	Claim is justified because estimated mean is less than 20 seconds	A2																																		
	(d) (i)	No	A1																																		
	(ii)	Any unsatisfied property, e.g. • distribution not symmetrical • histogram has a positive skew	A1	10																																	
10	(a)	Mean = $\frac{3 + 8 + 9 + 5 + 12 + 4 + 2 + 1}{8}$	M1																																		
		= 5.5 times	A1																																		
	(b)	All the numbers have the same frequency so there is no mode.	A1																																		
	(c)	Decrease	A1																																		
	(d)	People living closer to the cinema go to it more often.	A1																																		
	(e)	Quantitative	A1																																		
		Discrete	A1																																		
	(f)	Negative correlation, so people living closer to the cinema go more often.	A2	9																																	

11 (a) Statement B

A1


(b) % of people who prefer Running = $\frac{126}{360} \times 100$
= 35%

M1 MA1

A1

(c)

No. of Times	No. of People	Angle
Never	17	51°
Once	72	216°
Two or more	31	93°



MA4

(d) Last row is open-ended

A1

9

Total

80